



## News Release

### **Puma Biotechnology Licensing Partner Pint Pharma Receives Marketing Authorization for NERLYNX® (neratinib) for Extended Adjuvant Treatment of Early Stage Hormone Receptor Positive, HER2-Positive Breast Cancer**

- *First regulatory approval of NERLYNX in Latin America.*
- *NERLYNX is the first approved therapy to reduce the risk of invasive disease recurrence in HER2-positive breast cancer patients following trastuzumab-based adjuvant treatment.*
- *Up to 25% of HER2-positive early stage breast cancer patients treated with a trastuzumab-based regimen experience a recurrence.*

**LOS ANGELES, Calif., Sept. 4, 2019** – Puma Biotechnology, Inc. (Nasdaq: PBYI) announced that the Agency of Medicines, Food and Medical Devices (ANMAT) in Argentina has granted approval of NERLYNX® (neratinib) for the extended adjuvant treatment of adult patients with early stage hormone receptor positive HER2-overexpressed/amplified breast cancer and who completed adjuvant trastuzumab-based therapy less than one year ago.

ANMAT approval was based on the Phase III ExteNET trial, a multicenter, randomized, double-blind, placebo-controlled trial of neratinib following adjuvant trastuzumab treatment. Women (n=2,840) with early stage HER2-positive breast cancer and within two years of completing adjuvant trastuzumab therapy were randomized to receive either neratinib (n=1,420) or placebo (n=1,420) for one year.

The results of the ExteNET trial demonstrated that after two years of follow-up, for patients with hormone receptor positive, HER2-positive early stage breast cancer and who were treated within one year after the completion of trastuzumab based adjuvant therapy, invasive disease-free survival (iDFS) was 95.3% in the patients treated with neratinib compared with 90.8% in those receiving placebo (hazard ratio = 0.49; 95% CI: (0.30, 0.78); p=0.002).

The most common adverse reactions (>5%) were diarrhea, nausea, abdominal pain, fatigue, vomiting, rash, stomatitis, decreased appetite, muscle spasms, dyspepsia, AST or ALT increase, nail disorder, dry skin, abdominal distention, weight loss, and urinary tract infection. The most common adverse reaction leading to discontinuation was diarrhea, which was observed in 16.8% of neratinib-treated patients. Hepatotoxicity or increases in liver transaminases led to drug discontinuation in 1.7% of neratinib-treated patients.

Puma Biotechnology's Chairman, Chief Executive Officer and President Alan H. Auerbach, said, "We are pleased with the positive regulatory responses to NERLYNX that we have received internationally. The approval of NERLYNX in Argentina follows marketing authorizations in the United States, Europe, Australia and Canada. We look forward to collaborating with Pint Pharma and our other international partners to leverage their specific regulatory and commercial expertise as we make NERLYNX available to patients around the world."

#### **About HER2-Positive Breast Cancer**

Approximately 20 to 25 percent of breast cancer tumors over-express the HER2 protein and in the ExteNET study 57% of patients were found to have tumors that were hormone receptor positive. HER2-positive breast

cancer is often more aggressive than other types of breast cancer, increasing the risk of disease progression and death. Although research has shown that trastuzumab can reduce the risk of early stage HER2-positive breast cancer returning after surgery, up to 25% of patients treated with trastuzumab experience recurrence.

### **About Puma Biotechnology**

Puma Biotechnology, Inc. is a biopharmaceutical company with a focus on the development and commercialization of innovative products to enhance cancer care. Puma in-licenses the global development and commercialization rights to three drug candidates — PB272 (neratinib, oral), PB272 (neratinib, intravenous) and PB357. Neratinib, oral was approved by the U.S. Food and Drug Administration in July 2017 for the extended adjuvant treatment of adult patients with early stage HER2-overexpressed/amplified breast cancer, following adjuvant trastuzumab-based therapy, and is marketed in the United States as NERLYNX<sup>®</sup> (neratinib) tablets. NERLYNX was granted marketing authorization by the European Commission in September 2018 for the extended adjuvant treatment of adult patients with early stage hormone receptor-positive HER2-overexpressed/amplified breast cancer and who are less than one year from completion of prior adjuvant trastuzumab-based therapy. NERLYNX is a registered trademark of Puma Biotechnology, Inc.

Further information about Puma Biotechnology may be found at [www.pumabiotechnology.com](http://www.pumabiotechnology.com).

### **Important EU NERLYNX<sup>®</sup> (neratinib) Safety Information**

**All suspected adverse reactions should be reported in accordance with the national reporting system.**

The adverse reactions described in this section were identified in the randomized Phase 3 clinical trial (n=2840). The most common adverse reactions of any grade were diarrhoea (93.6%), nausea (42.5%), fatigue (27.3%), vomiting (26.8%), abdominal pain (22.7%), rash (15.4%), decreased appetite (13.7%), abdominal pain upper (13.2%), stomatitis (11.2%), and muscle spasms (10.0%).

The most common Grade 3-4 adverse reactions were diarrhoea (Grade 3, 36.9% and Grade 4, 0.2%) and vomiting (Grade 3, 3.4% and Grade 4, 0.1%).

Adverse reactions reported as serious included diarrhoea (1.9%), vomiting (1.3%), dehydration (1.1%), nausea (0.5%), alanine aminotransferase increased (0.4%), aspartate aminotransferase increased (0.4%), abdominal pain (0.3%), fatigue (0.3%) and decreased appetite (0.2%).

**For full European prescribing information, please refer to the NERLYNX (neratinib) Summary of Product Characteristics on the European Medicines Agency website (<http://www.ema.europa.eu/ema/>).**

### **Important Safety Information Regarding NERLYNX<sup>®</sup> (neratinib) U.S. Indication**

**NERLYNX<sup>®</sup> (neratinib) tablets, for oral use**

**INDICATIONS AND USAGE:** NERLYNX is a kinase inhibitor indicated for the extended adjuvant treatment of adult patients with HER2 overexpressed/amplified breast cancer, to follow adjuvant trastuzumab-based therapy.

**CONTRAINDICATIONS:** None

## WARNINGS AND PRECAUTIONS:

- **Diarrhea:** Aggressively manage diarrhea occurring despite recommended prophylaxis with additional antidiarrheals, fluids, and electrolytes as clinically indicated. Withhold NERLYNX in patients experiencing severe and/or persistent diarrhea. Permanently discontinue NERLYNX in patients experiencing Grade 4 diarrhea or Grade  $\geq 2$  diarrhea that occurs after maximal dose reduction.
- **Hepatotoxicity:** Monitor liver function tests monthly for the first 3 months of treatment, then every 3 months while on treatment and as clinically indicated. Withhold NERLYNX in patients experiencing Grade 3 liver abnormalities and permanently discontinue NERLYNX in patients experiencing Grade 4 liver abnormalities.
- **Embryo-Fetal Toxicity:** NERLYNX can cause fetal harm. Advise patients of potential risk to a fetus and to use effective contraception.

**ADVERSE REACTIONS:** The most common adverse reactions ( $\geq 5\%$ ) were diarrhea, nausea, abdominal pain, fatigue, vomiting, rash, stomatitis, decreased appetite, muscle spasms, dyspepsia, AST or ALT increase, nail disorder, dry skin, abdominal distention, weight decreased and urinary tract infection.

**To report SUSPECTED ADVERSE REACTIONS, contact Puma Biotechnology, Inc. at 1-844-NERLYNX (1-844-637-5969) and [www.NERLYNX.com](http://www.NERLYNX.com) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## DRUG INTERACTIONS:

- Gastric acid reducing agents: Avoid concomitant use with proton pump inhibitors. When patients require gastric acid reducing agents, use an H<sub>2</sub>-receptor antagonist or antacid. Separate NERLYNX by at least 3 hours with antacids. Separate NERLYNX by at least 2 hours before or 10 hours after H<sub>2</sub>-receptor antagonists.
- Strong or moderate CYP3A4 inhibitors: Avoid concomitant use.
- Strong or moderate CYP3A4 inducers: Avoid concomitant use.
- P-glycoprotein (P-gp) substrates: Monitor for adverse reactions of narrow therapeutic agents that are P-gp substrates when used concomitantly with NERLYNX.

## USE IN SPECIFIC POPULATIONS:

- **Lactation:** Advise women not to breastfeed.

Please see [Full Prescribing Information](#) for additional safety information.

The recommended dose of NERLYNX is 240 mg (six 40 mg tablets) given orally once daily with food, continuously for one year. Antidiarrheal prophylaxis should be initiated with the first dose of NERLYNX and continued during the first 2 months (56 days) of treatment and as needed thereafter.

To help ensure patients have access to NERLYNX, Puma has implemented the Puma Patient Lynx support program to assist patients and healthcare providers with reimbursement support and referrals to resources that can help with financial assistance. More information on the Puma Patient Lynx program can be found at [www.NERLYNX.com](http://www.NERLYNX.com) or 1-855-816-5421.

## **Forward-Looking Statements**

This press release contains forward-looking statements, including statements regarding the worldwide expansion of NERLYNX. All forward-looking statements involve risks and uncertainties that could cause the Company's actual results to differ materially from the anticipated results and expectations expressed in these forward-looking statements. These statements are based on current expectations, forecasts and assumptions, and actual outcomes and results could differ materially from these statements due to a number of factors, which include, but are not limited to, the risk factors disclosed in the periodic and current reports filed by the Company with the Securities and Exchange Commission from time to time, including the Company's Annual Report on Form 10-K for the year ended December 31, 2018. Readers are cautioned not to place undue reliance on these forward-looking statements, which speak only as of the date hereof. The Company assumes no obligation to update these forward-looking statements, except as required by law.

## **Contact**

Alan H. Auerbach or Mariann Ohanesian, Puma Biotechnology, Inc., +1 424 248 6500  
info@pumabiotechnology.com  
ir@pumabiotechnology.com

David Schull or Maggie Beller, Russo Partners, +1-212-845-4200  
david.schull@russopartnersllc.com  
maggie.beller@russopartnersllc.com